



University of  
Zurich<sup>UZH</sup>

Zurich Open Repository and  
Archive

University of Zurich  
Main Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2006

---

## Chemoselectivity of the [2+3]-Cycloaddition of Thiocarbonyl Ylides with 5-Benzylidene-3-phenylrhodanine

Seyfried, M S ; Linden, Anthony ; Młostoń, Grzegorz ; Heimgartner, H

**Abstract:** Reactions of three different thiocarbonyl S-methylides, generated from thiobenzophenone (2), 2,2,4,4-tetramethyl-3-thioxocyclobutanone (3), and adamantanethione (8), respectively, and diazomethane, with 5-benzylidene-3-phenylrhodanine (12) were carried out. The aromatic thiocarbonyl ylide 1a adds chemoselectively to the C,C-double bond, but the spirocyclic 1,3-dithiolane 18, i.e. the [2+3]-cycloadduct with the C=S group of 12, was also formed as a minor product. In the cases of the aliphatic thiocarbonyl ylides 6 and 20, the [2+3]-cycloaddition occurred at the exocyclic C,C-double bond exclusively to give the spirocyclic tetrahydrothiophene derivatives 23 and 21, respectively. A smooth acid-catalyzed decomposition of 18 yielded the 2-diphenylmethylidene derivative 19. The formation of product 24, which was obtained in the reaction of the sterically congested ylide 6 with 12, is explained by a 1,4-H-shift in an intermediate zwitterionic adduct. The structures of the tetrahydrothiophenes 17, 21 and 23, as well as that of 24, were established by X-ray crystallography.

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-54086>

Journal Article

Published Version

Originally published at:

Seyfried, M S; Linden, Anthony; Młostoń, Grzegorz; Heimgartner, H (2006). Chemoselectivity of the [2+3]-Cycloaddition of Thiocarbonyl Ylides with 5-Benzylidene-3-phenylrhodanine. *Polish Journal of Chemistry*, 80(8):1363-1376.

## Chemoselectivity of the [2+3]-Cycloaddition of Thiocarbonyl Ylides with 5-Benzylidene-3-phenylrhodanine

by M.S. Seyfried<sup>1\*</sup>, A. Linden<sup>1</sup>, G. Mlostoń<sup>2</sup> and H. Heimgartner<sup>1\*\*</sup>

<sup>1</sup>*Institute of Organic Chemistry, University of Zürich,  
Winterthurerstrasse 190, CH-8057 Zürich, Switzerland*

<sup>2</sup>*Section of Heteroorganic Compounds, University of Łódź, Narutowicza 68, PL-90-136 Łódź, Poland*

(Received March 27th, 2006)

Reactions of three different thiocarbonyl *S*-methylides, generated from thiobenzophenone (**2**), 2,2,4,4-tetramethyl-3-thioxocyclobutanone (**3**), and adamantanethione (**8**), respectively, and diazomethane, with 5-benzylidene-3-phenylrhodanine (**12**) were carried out. The aromatic thiocarbonyl ylide **1a** adds chemoselectively to the C,C-double bond, but the spirocyclic 1,3-dithiolane **18**, *i.e.* the [2+3]-cycloadduct with the C=S group of **12**, was also formed as a minor product. In the cases of the aliphatic thiocarbonyl ylides **6** and **20**, the [2+3]-cycloaddition occurred at the exocyclic C,C-double bond exclusively to give the spirocyclic tetrahydrothiophene derivatives **23** and **21**, respectively. A smooth acid-catalyzed decomposition of **18** yielded the 2-diphenylmethylidene derivative **19**. The formation of product **24**, which was obtained in the reaction of the sterically congested ylide **6** with **12**, is explained by a 1,4-H-shift in an intermediate zwitterionic adduct. The structures of the tetrahydrothiophenes **17**, **21** and **23**, as well as that of **24**, were established by X-ray crystallography.

**Key words:** thiocarbonyl ylides, 1,3-dipolar cycloaddition, rhodanine, 2,5-dihydro-1,3,4-thiadiazoles, crystal structure

Although the fruitful concept of 1,3-dipolar cycloaddition was formulated by Huisgen as early as 1963 [1] (see also [2,3]), and some [2+3] cycloadditions with thioketones were observed early in the last century [4,5],<sup>\*\*\*</sup> the usefulness of the C=S group as a dipolarophile was recognized only in the 1980s [6,7]. Since then, a large series of [2+3] cycloadditions with C=S compounds have been investigated and their reaction mechanisms established. As a result of kinetic studies, thioketones are now recognized as extraordinarily reactive dipolarophiles and are called ‘superdipolarophiles’ [8–11].

In the last few years, interest in thiocarbonyl ylides as 1,3-dipoles has increased [12,13]. Their reaction with C=S containing compounds, such as aromatic and cycloaliphatic thioketones [14–16], dithioesters [15,16], trithiocarbonates [15], and 1,3-thiazole-5(4*H*)-thiones [17], leads to 1,3-dithiolanes (‘Schönberg reaction’ [14,18],


\* Part of the diploma thesis of M.S.S., University of Zürich, 2005.

\*\* Author for correspondence; Prof. H. Heimgartner, e-mail: heimgart@oci.unizh.ch

\*\*\* At that time, the mechanism of the formation of the described products was unknown.

Scheme 1). Aromatic thioketone *S*-methylides **1** react with aromatic as well as cycloaliphatic thioketones (*e.g.* **2** and **3**) to give the sterically crowded 1,3-dithiolanes of type **4** and **5**, respectively, in a regioselective manner [14,15,19]. On the other hand, the reactions of cycloaliphatic thioketone *S*-methylides, such as **6** with **2**, lead to mixtures of 1,3-dithiolanes, *e.g.* **5** and **7** in favor of the sterically congested **5** [16,18,20]. Finally, the sterically less crowded products of type **9** are formed regioselectively in reactions of cycloaliphatic thioketone *S*-methylides with aliphatic thioketones (*e.g.* **6** and **8**) [16,18,20].


Scheme 1



In addition to thioketones, heterocyclic thiones **10–12** (Scheme 2) have been used as C=S dipolarophiles. For example, benzonitrile oxides, imides and ylides [21], diazo compounds [22], azomethine ylides [23], carbonyl ylides [24] and thiocarbonyl ylides [17] react exclusively with the C=S group of **10**. On the other hand, benzonitrile oxide (**15**) and nitrones add chemoselectively to the C,C-bond of **11** [25], whereas reactions of benzonitrile imides **13** and benzonitrile oxide (**15**) with the rhodanine derivative **12** yield cycloadducts **14** [26] and 1,3-thiazolidine-2,4-dione **16**, respectively. In the latter case, the initially formed cycloadduct, a spirocyclic 1,4,2-oxathiazole derivative, is not stable and a subsequent cycloreversion leads to the final product [27].

Recently, we investigated further 1,3-dipolar cycloadditions with **12**, which contains three different  $\pi$ -systems as potential dipolarophilic groups. In the present paper, new results of the reactions of **12** with thiocarbonyl *S*-methylides **1a**, **6** and **20** are discussed.


Scheme 2




## RESULTS AND DISCUSSION

According to the protocol in [15], thiobenzophenone *S*-methylide (**1a**) was generated *in situ* in dichloromethane at low temperature in the presence of the rhodanine derivative **12** [28]. After stirring overnight at room temperature, chromatographic workup gave two 1:1 adducts **17** and **18**, which were isolated in 61 and 3% yield, respectively (Scheme 3). Furthermore, 30% of the starting material **12** was recovered. The ESI-MS of the two adducts showed that they are isomers ( $m/z = 532$  ( $100, [M + \text{Na}]^+$ )), and the  $^{13}\text{C}$ -NMR spectra revealed the presence of a  $\text{C}=\text{S}$  group in **17** (198.9 ppm) and the absence of this group in **18**. As in both cases a carbonyl group was detected (IR,  $^{13}\text{C}$ -NMR), **17** is the  $\text{C}=\text{C}$  adduct and **18** the expected  $\text{C}=\text{S}$  adduct. The structure of **17** was established by an X-ray crystal structure analysis (Figure 1).

Scheme 3






**Figure 1.** ORTEP-plot [29] of the molecular structure of **17** (arbitrary numbering of the atoms; 50% probability ellipsoids).

In the case of **18**, the structure was assigned on the basis of the spectroscopic data. The regioselectivity of the cycloaddition, *i.e.* the formation of the more crowded 1,3-dithiolane, follows from the NMR absorption of the CH<sub>2</sub> group (*AB* at 3.84 and 3.59 ppm, *J* = 8.4 Hz, *t* at 31.9 ppm; see [15,18]). Under slightly acidic conditions in CDCl<sub>3</sub>, **18** decomposed to give the known product **19**<sup>\*</sup> in quantitative yield. The structure of product **19** proves the presence of a C,C bond between Ph<sub>2</sub>C and C(2) of the 1,3-thiazolidine ring in **18**. A reaction mechanism for the decomposition of **18** is proposed in Scheme 4 (see also [31]). Elimination of thioformaldehyde under acid catalysis could lead to intermediate **A**, which forms the spirocyclic thiirane **B** *via* nucleophilic attack of the thiole group at the iminium C-atom. Subsequently, elimination of sulfur from **B** yields the isolated product **19**. The last reaction is also responsible for the formation of **19** in the reaction of **12** with diphenyldiazomethane [30].

Similarly, the reaction of **12** with adamantanethione *S*-methylide (**20**) was carried out: diazomethane was added to a solution of adamantanethione (**8**) in toluene at −20°C and, after decolorization of the solution, **12** was added and the mixture


<sup>\*</sup>Compound **19** has been prepared independently of **12** and diphenyldiazomethane in refluxing toluene [30].

Scheme 4




refluxed for 12 h. Chromatographic workup gave 28% of the 1:1-cycloadduct **21** and 29% of the starting material **12** (Scheme 5). No additional product could be detected. The relatively low yield of **21** can be explained by the fact that the treatment of **8** with diazomethane leads to a mixture of the 1,3,4-thiadiazole (the precursor of **20**) and the 1,2,3-isomer [16]. The latter does not eliminate  $\text{N}_2$  to give **20** on heating but decomposes during chromatographic workup. The structure of **21** was established by X-ray crystallography (Figure 2). These results show that the 1,3-dipolar cycloaddition of **20** with **12** occurred exclusively at the C,C-double bond, and not at the thiocarbonyl group. Furthermore, it is worth mentioning that the sterically less favored cycloadduct is formed.

Scheme 5




Finally, the reaction of 2,2,4,4-tetramethyl-3-oxocyclobutanethione *S*-methylide (**6**) with **12** was performed. The precursor of **6**, the 2,5-dihydro-1,3,4-thiadiazole **22**, was prepared from 2,2,4,4-tetramethyl-3-thioxobutanone (**3**) and diazomethane in diethyl ether at  $-78^\circ\text{C}$  and crystallized from pentane [20]. A solution of **12** and 1.4 equivalent of **22** in THF was heated to  $50^\circ\text{C}$ . The nitrogen evolution ceased after 2 h. Then, the solvent was evaporated and the residue separated by means of column chromatography. In addition to 74% of the starting material **12**, two isomeric products




**Figure 2.** ORTEP-plot [29] of the molecular structure of **21** (arbitrary numbering of the atoms; 50% probability ellipsoids).

with the molecular weight of 1:1-adducts (ESI-MS:  $m/z = 490$  ( $100$ ,  $[M + Na]^+$ )) were obtained in **8** and 10% yield, respectively\*. Based on the spectroscopic data, the first product was assigned the structure of the cycloadduct **23** (Scheme 6). The regioselectivity of the cycloaddition was indicated by the similarity of the NMR absorptions of the  $-\text{CH}_2-\text{CHPh}-$  fragment in adducts **17**, **21** and **23**. Finally, the structure of **23** was established by X-ray crystallography (Figure 3).

Scheme 6



\* In the crude reaction mixture, significant amounts of 4,4,6,6-tetramethyl-1-thiaspiro[2.3]hexan-5-one [32], the product of the 1,3-dipolar electrocycloaddition of the thiocarbonyl ylide **6**, were detected.




**Figure 3.** ORTEP-plots [29] of the molecular structures of one of the two symmetry-independent molecules of **23** (top) and of **24** (arbitrary numbering of the atoms; 50% probability ellipsoids).

The surprising structure of the second product, **24**, was again established by an X-ray crystal structure determination (Figure 3). A likely explanation for its formation is presented in Scheme 7 and is based on the assumption that the reaction with **6** may occur stepwise, in analogy to earlier described reactions of **6** with electron-deficient ethenes [13,33]. Addition of the nucleophilic thiocarbonyl ylide **6** to the exocyclic C,C-double bond of **12** in a Michael-type reaction could lead to the intermediate **C**, which gives the isolated products **23** and **24** via a 1,5-dipolar electrocyclization and a 1,4-H-shift, respectively.



Scheme 7



In conclusion, the experiments show that thiocarbonyl *S*-methanides **1a**, **6** and **20** react with **12** chemoselectively at the C,C-double bond to give spirocyclic tetrahydrothiophene derivatives. A 1,3-dithiolane as a minor adduct was observed only in the case of thiobenzophenone *S*-methylide (**1a**) and results from the addition occurring at the C=S group. This is a surprising result as, in general, C=S groups are known to be superior dipolarophiles, and benzonitrile oxide and -imides react with **12** at the C=S group exclusively. A possible explanation of the observed results is the high nucleophilicity of thiocarbonyl ylides and, therefore, a preference for nucleophilic addition onto the  $\alpha,\beta$ -unsaturated ketone. In all examples, only one regioisomer, *i.e.* the sterically more crowded adduct was formed. Furthermore, the products consist of a single diastereoisomer as a racemate. The formation of **24** in the case of the sterically most congested thiocarbonyl *S*-methanide **6** is most likely a result of an H-shift in intermediate **C** (Scheme 7), which also could be the precursor of the cycloadduct **23** in a two-step reaction.

## EXPERIMENTAL

**1. General.** See [16]. The  $^{13}\text{C}$ -NMR spectra were recorded by using DEPT registration.

**2. Starting materials.** (Z)-5-Benzylidene-3-phenyl-2-thioxo-1,3-thiazolidin-4-one (benzylidene-phenylrhodanine, **12**) was prepared in two steps following the protocol described in [28] (see also [26b]): treatment of 2-mercaptoacetic acid with phenyl isothiocyanate in a mixture of  $\text{H}_2\text{O}$  and EtOH gave *N*-phenylrhodanine in 77% yield. The latter reacted with benzaldehyde in refluxing acetic acid/sodium acetate to give **12** in 85% yield. Thiobenzophenone (**2**) [34,35], 2,2,4,4-tetramethyl-3-thioxocyclobutanone (**3**) [36,37], and adamantanethione (**8**) [35,38] were synthesized according to known procedures.

**3. Reaction of 12 with thiobenzophenone *S*-methylide (1a).** A solution of **2** (0.612 g, 3.09 mmol) in  $\text{CH}_2\text{Cl}_2$  (150 ml) under an argon atmosphere was cooled to  $-78^\circ\text{C}$ , and 7.4 ml of a 0.5 M solution of diazomethane (*ca.* 3.7 mmol) in Et<sub>2</sub>O were added, whereby the blue color of the solution disappeared. Then, a solution of **2** (0.918 g, 3.09 mmol) in  $\text{CH}_2\text{Cl}_2$  was added and the solution was allowed to reach room temperature while stirring. After *ca.* 15 h, the solvent was evaporated and the residue was separated by column chromatography ( $\text{SiO}_2$ , hexane/ethyl acetate 5:1) to give **12** (0.271 g, 30%), **17** (0.964 g, 61%), and **18** (46 mg, 3%).

*cis*-3,6,6,9-Tetraphenyl-2-thioxo-3-aza-1,7-dithiaspiro[4.4]nonan-4-one (**17**): Colorless crystals, m.p.  $264.0\text{--}264.5^\circ\text{C}$  ( $\text{CH}_2\text{Cl}_2$ /hexane).  $R_f$  (hexane/ethyl acetate 5:1): 0.38. IR (KBr):  $3057_{\text{w}}$ ,  $2934_{\text{vw}}$ ,  $1956_{\text{vw}}$ ,  $1736_{\text{vs}}$ ,  $1681_{\text{w}}$ ,  $1593_{\text{w}}$ ,  $1492_{\text{s}}$ ,  $1454_{\text{m}}$ ,  $1444_{\text{m}}$ ,  $1343_{\text{s}}$ ,  $1291_{\text{w}}$ ,  $1243_{\text{vs}}$ ,  $1180_{\text{m}}$ ,  $1156_{\text{m}}$ ,  $1072_{\text{m}}$ ,  $1058_{\text{w}}$ ,  $1032_{\text{w}}$ ,  $1003_{\text{w}}$ ,  $779_{\text{w}}$ ,  $766_{\text{w}}$ ,  $740_{\text{s}}$ ,  $698_{\text{vs}}$ ,  $622_{\text{w}}$ ,  $613_{\text{w}}$ .  $^1\text{H}$ -NMR (300 MHz,  $\text{CDCl}_3$ ):  $7.57\text{--}7.54$

(*m*, 2 arom. H); 7.42–7.24 (*m*, 18 arom. H); 4.35 (*t*,  $J = 10.4$ , PhCH); 3.62 (*dd*-like,  $J = 8.2$ , 10.4, CH<sub>2</sub>). <sup>13</sup>C-NMR (75.5 MHz, CDCl<sub>3</sub>): 198.9 (*s*, CS); 172.4 (*s*, CO); 142.7, 140.6, 135.0, 133.3 (4*s*, 4 arom. C); 130.5, 130.0, 129.4, 129.2, 129.0, 128.4, 128.3, 127.9, 127.6, 127.5, 127.1, 126.8 (12*d*, 20 arom. CH); 80.4, 74.2 (2*s*, spiro-C, Ph<sub>2</sub>C); 55.9 (*d*, PhCH); 32.5 (*t*, CH<sub>2</sub>). ESI-MS (CH<sub>2</sub>Cl<sub>2</sub>/MeCN 1:1 + NaI): 532 (100, [M + Na]<sup>+</sup>). Anal. Calc. for C<sub>30</sub>H<sub>23</sub>NOS<sub>3</sub> (509.70): C 70.69, H 4.55, N 2.75, S 18.87. Found: C 70.64, H 4.54, N 2.70, S 18.86.

Suitable crystals for an X-ray crystal-structure determination were obtained by isothermal distillation of pentane into a solution of **17** in CH<sub>2</sub>Cl<sub>2</sub>/hexane.

(*Z*)-7-Benzylidene-4,4,9-triphenyl-9-aza-1,3,6-trithiaspiro[4.4]nonan-8-one (**18**). Colorless solid, m.p. 211–212°C (CH<sub>2</sub>Cl<sub>2</sub>/hexane). *R*<sub>f</sub> (hexane/ethyl acetate 5:1): 0.25. IR (KBr): 3056*w*, 2981*vw*, 2912*vw*, 1767*w*, 1684*vs*, 1608*w*, 1491*s*, 1444*m*, 1411*w*, 1332*vs*, 1286*w*, 1224*w*, 1189*s*, 1115*m*, 1078*w*, 1034*m*, 1019*m*, 918*w*, 884*w*, 825*w*, 785*w*, 761*w*, 749*w*, 719*s*, 695*s*, 642*w*, 607*m*. <sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>): 7.63–7.20 (*m*, 20 arom. H, PhCH); 3.84, 3.59 (*AB*,  $J = 8.4$ , CH<sub>2</sub>). <sup>13</sup>C-NMR (75.5 MHz, CDCl<sub>3</sub>): 166.8 (*s*, CO); 144.3, 139.6, 138.6, 134.6 (4*s*, 4 arom. C); 129.7, 129.5, 129.0, 128.8, 128.3, 127.4 (6*d*, 20 arom. CH, PhCH); 123.8 (*s*, PhCH=C); 93.4 (*s*, spiro-C); 76.3 (*s*, Ph<sub>2</sub>C); 31.9 (*t*, SCH<sub>2</sub>S). ESI-MS (MeOH + NaI): 532 (100, [M + Na]<sup>+</sup>).

(*Z*)-5-Benzylidene-2-diphenylmethylene-3-phenyl-1,3-thiazolidin-4-one (**19**). A slightly acidic solution of **18** in CDCl<sub>3</sub> turned yellow and a solid precipitated. Filtration and evaporation of the solid gave **19** in quantitative yield.\*

**4. Reaction of 12 with adamantanethione S-methylide (20).** A solution of **8** (0.165 g, 0.99 mmol) in toluene (50 ml) under an argon atmosphere was cooled to –20°C, and 2.6 ml of a 0.5 M solution of diazomethane (*ca.* 1.3 mmol) were added in two portions. After stirring for 2 h, the orange color of the solution vanished, and a solution of **12** (0.298 g, 1.00 mmol) in toluene (10 ml) was added. The mixture was heated to reflux (*ca.* 140°C) for 12 h, then the solvent was evaporated and the residue was separated by column chromatography (SiO<sub>2</sub>, hexane/ethyl acetate 5:1) to yield **12** (0.206 g, 69%) and **21** (0.125 g, 28%).

Spiro[4-oxo-3,9-diphenyl-2-thioxo-3-aza-1,7-dithiaspiro[4.4]nonane-6,2'-tricyclo[3.3.1.1<sup>3,7</sup>]decane] (**21**). Pale yellow crystals, m.p. 211–213°C (CH<sub>2</sub>Cl<sub>2</sub>/hexane). *R*<sub>f</sub> (hexane/ethyl acetate 5:1): 0.55. IR (KBr): 3057*w*, 3031*vw*, 2937*m*, 2908*vs*, 2867*m*, 2851*s*, 1726*vs*, 1593*w*, 1493*m*, 1472*w*, 1453*m*, 1446*m*, 1340*s*, 1314*w*, 1291*w*, 1241*vs*, 1176*m*, 1166*m*, 1138*w*, 1098*m*, 1084*w*, 1072*w*, 1026*w*, 991*w*, 950*w*, 913*w*, 839*w*, 770*w*, 751*w*, 738*m*, 730*m*, 699*s*, 689*s*, 611*w*. <sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>): 7.48–7.31 (*m*, 10 arom. H); 4.51 (*dd*,  $J(\text{trans}) = 11.7$ ,  $J(\text{cis}) = 7.3$ , PhCH); 3.28–3.12 (*m*, CH<sub>2</sub>); 2.75 (*br. s*, 1 H); 2.68–2.65 (*m*, 2 H); 2.48 (*br. d*,  $J = 13.9$ , 1 H); 2.14 (*br. d*,  $J = 13.0$ , 1 H); 1.95–1.59 (*m*, 9 H). <sup>13</sup>C-NMR (75.5 MHz, CDCl<sub>3</sub>): 199.0 (*s*, CS); 175.0 (*s*, CO); 135.2, 134.5 (2*s*, 2 arom. C); 130.3, 129.4, 129.3, 128.6, 128.1, 127.8 (6*d*, 10 arom. CH); 81.7, 73.9 (2*s*, 2 spiro-C); 58.3 (*d*, PhCH); 39.6 (*d*, adamantane CH); 38.0 (*t*, 2 adamantane CH<sub>2</sub>); 37.4 (*d*, adamantane CH); 36.7, 35.8, 34.1, 31.7 (4*t*, 4 adamantane CH<sub>2</sub>); 26.7, 25.9 (2*d*, 2 adamantane CH). ESI-MS (CH<sub>2</sub>Cl<sub>2</sub>/MeCN + NaI): 500 (100, [M + Na]<sup>+</sup>). Anal. Calc. for C<sub>27</sub>H<sub>27</sub>NOS<sub>3</sub> (477.70): C 67.88, H 5.70, N 2.93, S 20.14. Found: C 67.98, H 5.62, N 2.84, S 20.15.

Suitable crystals for an X-ray crystal-structure determination of **21** were obtained by isothermal distillation of pentane into a solution of **21** in CH<sub>2</sub>Cl<sub>2</sub>/hexane.

**5. Reaction of 12 with 2,2,4,4-tetramethyl-3-thioxocyclobutane S-methylide (6).** A solution of **3** (0.628 g, 4.02 mmol) in Et<sub>2</sub>O (10 ml) under an argon atmosphere was cooled to –78°C, and 15 ml of a 0.5 M solution of diazomethane (*ca.* 7.5 mmol) were added in two portions. The red color of the solution disappeared after stirring for 2 h. Then, the solvent was removed at 0°C by bubbling N<sub>2</sub> gas through the solution. The solid residue was dissolved with pentane (15 ml) at 0°C, the solution was cooled to –78°C, and the crystallized *1,1,3,3-tetramethyl-5-thia-7,8-diazaspiro[3.4]oct-7-en-2-one* (**22**) was isolated by decantation. Yield: 0.694 g (87%). A solution of **22** (0.596 g, 3.00 mmol) and **12** (0.638 g, 2.14 mmol) in THF (10 ml) under an argon atmosphere was heated to 50°C, when the evolution of N<sub>2</sub> started. After 2 h, the evolution of N<sub>2</sub> ceased (64 ml N<sub>2</sub>, 2.9 mmol), the solvent was evaporated and the residue was separated by column chromatography (SiO<sub>2</sub>, hexane/ethyl acetate 5:1) to give **12** (467 mg, 74%), **23** (81 mg, 8%), and **24** (104 mg, 10%).

\*The same product was obtained in 93% yield from **12** and diphenyldiazomethane in toluene [30].

*1,1,3,3-Tetramethyl-8,10-diphenyl-7-thioxo-8-aza-6,12-dithiaspiro[3.0.4.3]dodecane-2,9-dione (23).* Pale yellow crystals, m.p. 208–209°C.  $R_f$  (hexane/ethyl acetate 5:1): 0.34. IR (KBr): 3054vw, 2968w, 1783vs, 1738vs, 1680m, 1494s, 1471m, 1455m, 1417m, 1383m, 1352s, 1246vs, 1176s, 1135m, 1090m, 971m, 900m, 844m, 825m, 779m, 724m, 700s, 687m, 613m.  $^1\text{H-NMR}$  (300 MHz,  $\text{CDCl}_3$ ): 7.56–7.27 (m, 10 arom. H); 4.13 (t,  $J = 9.6$ , PhCH); 3.40–3.31 (m,  $\text{CH}_2$ ); 1.82, 1.66, 1.65, 1.39 (4s, 4 Me).  $^{13}\text{C-NMR}$  (75.5 MHz,  $\text{CDCl}_3$ ): 217.3 (s, CO); 199.2 (s, CS); 172.5 (s, CO); 135.2, 133.7 (2s, 2 arom. C); 129.7, 129.5, 129.3, 129.1, 128.9, 128.5, 128.3, 127.6 (8d, 10 arom. CH); 82.5, 74.3 (2s, 2 spiro-C); 69.7, 62.7 (2s, 2 Me<sub>2</sub>C); 57.81 (d, PhCH); 32.1 (t,  $\text{CH}_2$ ); 25.7, 24.3, 22.5, 22.1 (4q, 4 Me). ESI-MS ( $\text{MeCN}/\text{CH}_2\text{Cl}_2$  1:1 + NaI): 490 (100,  $[\text{M} + \text{Na}]^+$ ). Anal. Calc. for  $\text{C}_{25}\text{H}_{25}\text{NO}_2\text{S}_3$  (467.67): C 64.21, H 5.39, N 3.00, S 20.57. Found: C 64.02, H 5.36, N 2.87, S 20.50.

Suitable crystals for an X-ray crystal-structure determination were obtained by isothermal distillation of pentane into a solution of **23** in 1,2-dimethoxyethane.

*3-Phenyl-5-{1-phenyl-2-[(2',2',4',4'-tetramethylcyclobutan-1'-yl)thio]ethylidene}-2-thioxo-1,3-thiazolidine-3',4'-dione (24).* Orange crystals, m.p. 154–158°C.  $R_f$  (hexane/ethyl acetate 5:1): 0.31. IR (KBr): 3045vw, 2959m, 2923w, 2861w, 1770vs, 1710vs, 1590m, 1569w, 1491m, 1456m, 1441w, 1415w, 1379w, 1364w, 1347s, 1315w, 1291w, 1247vs, 1183s, 1157m, 1092w, 1073w, 1024m, 914w, 841w, 764w, 729m, 700m, 689m, 639w, 630w, 612w.  $^1\text{H-NMR}$  (300 MHz,  $\text{CDCl}_3$ ): 7.58–7.43 (m, 7 arom. H); 7.38 (br. s, 1 arom. H); 7.26–7.23 (m, 2 arom. H); 4.38 (s,  $\text{CH}_2$ ); 3.09 (s, CHS); 1.15, 1.14 (2s, 4 Me).  $^{13}\text{C-NMR}$  (75.5 MHz,  $\text{CDCl}_3$ ): 219.9 (s, CO); 193.5 (s, CS); 165.4 (s, CO); 149.4 (s, PhC=C); 139.6, 134.9 (2s, 2 arom. C); 130.1, 129.6, 129.5, 129.2, 128.3, 128.0, 127.2 (7d, 10 arom. CH); 125.3 (s, PhC=C); 60.3 (s, 2 Me<sub>2</sub>C); 55.6 (d, CHS); 34.3 (t,  $\text{CH}_2$ ); 24.4, 19.8 (2q, 4 Me). ESI-MS ( $\text{CH}_2\text{Cl}_2$  + NaI): 490 (100,  $[\text{M} + \text{Na}]^+$ ), 468 (18), 420 (31), 333 (9), 304 (16).

Suitable crystals for an X-ray crystal-structure determination of **24** were obtained by isothermal distillation of pentane into a solution of **24** in  $\text{CDCl}_3$ /hexane.

**6. X-ray crystal-structure determination of 17, 21, 23 and 24** (see Table 1 and Figs. 1–3)\*. All measurements were made on a Nonius KappaCCD diffractometer [39] using graphite-monochromated  $\text{MoK}\alpha$  radiation ( $\lambda$  0.71073 Å) and an Oxford Cryosystems Cryostream 700 cooler. Data reduction was performed with HKL Denzo and Scalepack [40]. The intensities were corrected for Lorentz and polarization effects, and absorption corrections based on the multi-scan method [41] were applied. Equivalent reflections were merged. The data collection and refinement parameters are given in Table 1, and views of the molecules are shown in Figs. 1–3. The structures of **17**, **21** and **23** were solved by direct methods using SIR92 [42], which revealed the positions of all non-H-atoms. In the case of **23**, there are two symmetry-independent molecules in the asymmetric unit. The coordinates of the two molecules were tested carefully for a relationship from a higher symmetry space group using the program PLATON [43], but none could be found. The structure of **24** was solved by heavy-atom Patterson methods [44], which revealed the positions of the S-atoms. All remaining non-H-atoms were located in a Fourier expansion of the Patterson solution, which was performed by DIRDIF94 [45]. For each structure, the non-H-atoms were refined anisotropically. All of the H-atoms were placed in geometrically calculated positions and refined using a riding model where each H-atom was assigned a fixed isotropic displacement parameter with a value equal to  $1.2U_{\text{eq}}$  of its parent C-atom ( $1.5U_{\text{eq}}$  for the Me groups of **23** and **24**). The refinement of each structure was carried out on  $F^2$  using full-matrix least-squares procedures, which minimized the function  $\sum w(F_o^2 - F_c^2)^2$ . A correction for secondary extinction was applied for **23**. In the cases of **17**, **21** and **23**, three, two and eight reflections, respectively, whose intensities were considered to be extreme outliers, were omitted from the final refinement. For **23**, the largest peak of residual electron density was 2.36 Å from atom H(15), but could not be rationalised with a chemically logical position, nor could it be refined successfully as a partial occupancy water molecule. Neutral atom scattering factors for non-H-atoms were taken from [46a], and the scattering factors for H-atoms were taken from [47]. Anomalous dispersion effects were included in  $F_c$  [48]; the values for  $f'$  and  $f''$  were those of [46b]. The values of the mass attenuation coefficients are those of [46c]. All calculations were performed using the SHELXL97 [49] program.

\*CCDC-294731–294734 contain the supplementary crystallographic data for this paper.

These data can be obtained free of charge from the Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).

**Table 1.** Crystallographic data for compounds **17**, **21**, **23** and **24**.

	<b>17</b>	<b>21</b>
Crystallized from	CH <sub>2</sub> Cl <sub>2</sub> /hexane/pentane	CH <sub>2</sub> Cl <sub>2</sub> /hexane/pentane
Empirical formula	C <sub>30</sub> H <sub>23</sub> NOS <sub>3</sub>	C <sub>27</sub> H <sub>27</sub> NOS <sub>3</sub>
Formula weight [g mol <sup>-1</sup> ]	509.70	477.70
Crystal color, habit	colorless, prism	pale yellow, prism
Crystal dimensions [mm]	0.15 × 0.20 × 0.25	0.17 × 0.20 × 0.30
Temperature [K]	160(1)	160(1)
Crystal system	triclinic	monoclinic
Space group	<i>P</i> $\bar{1}$	<i>P</i> 2 <sub>1</sub> / <i>n</i>
<i>Z</i>	2	4
Reflections for cell determination	46235	34682
2 $\theta$ range for cell determination [°]	4–60	4–60
Unit cell parameters		
<i>a</i> [Å]	10.3260(2)	12.7901(2)
<i>b</i> [Å]	11.2042(2)	10.9689(2)
<i>c</i> [Å]	11.3691(1)	16.3241(2)
$\alpha$ [°]	98.8038(9)	90
$\beta$ [°]	92.4303(9)	101.2457(8)
$\gamma$ [°]	107.7516(7)	90
<i>V</i> [Å <sup>3</sup> ]	1232.40(3)	2246.19(6)
<i>D</i> <sub>x</sub> [g cm <sup>-3</sup> ]	1.373	1.412
$\mu$ (MoK $\alpha$ ) [mm <sup>-1</sup> ]	0.326	0.352
Scan type	$\phi$ and $\omega$	$\phi$ and $\omega$
2 $\theta$ (max) [°]	60	60
Transmission factors (min; max)	0.871; 0.955	0.864; 0.942
Total reflections measured	36407	60592
Symmetry independent reflections	7170	6570
Reflections with <i>I</i> > 2 $\sigma$ ( <i>I</i> )	5809	5505
Reflections used in refinement	7167	6568
Parameters refined	316	289
Final <i>R</i> ( <i>F</i> ) [ <i>I</i> > 2 $\sigma$ ( <i>I</i> ) reflections]	0.0386	0.0348
<i>wR</i> ( <i>F</i> <sup>2</sup> ) (all data)	0.1034	0.0923
Weights: <sup>a)</sup> <i>a</i> ; <i>b</i>	0.05; 0.509	0.0441; 0.9598
Goodness of fit	1.024	1.053
Secondary extinction coefficient	—	—
Final $\Delta$ <sub>max</sub> / $\sigma$	0.001	0.001
$\Delta\rho$ (max; min) [e Å <sup>-3</sup> ]	0.34; -0.37	0.35; -0.36

<sup>a)</sup>  $w = [\sigma^2(F_o^2) + (aP)^2 + bP]^{-1}$ , where  $P = (F_o^2 + 2F_c^2) / 3$

**Table 1.** Crystallographic data for compounds **17**, **21**, **23** and **24** (continued).

	<b>23</b>	<b>24</b>
Crystallized from	DMOE/pentane	CDCl <sub>3</sub> /hexane/pentane
Empirical formula	C <sub>25</sub> H <sub>25</sub> NO <sub>2</sub> S <sub>3</sub>	C <sub>25</sub> H <sub>25</sub> NO <sub>2</sub> S <sub>3</sub>
Formula weight [g mol <sup>-1</sup> ]	467.66	467.66
Crystal color, habit	pale yellow, prism	orange, prism
Crystal dimensions [mm]	0.20 × 0.25 × 0.28	0.22 × 0.25 × 0.28
Temperature [K]	160(1)	160(1)
Crystal system	triclinic	triclinic
Space group	<i>P</i> $\bar{1}$	<i>P</i> $\bar{1}$
<i>Z</i>	4	2
Reflections for cell determination	207466	55709
2 $\theta$ range for cell determination [°]	4–55	4–60
Unit cell parameters		
<i>a</i> [Å]	8.4135(2)	8.0529(2)
<i>b</i> [Å]	15.7231(4)	11.4487(2)
<i>c</i> [Å]	19.1282(5)	13.2158(2)
$\alpha$ [°]	111.005(1)	79.4247(9)
$\beta$ [°]	90.423(2)	85.547(1)
$\gamma$ [°]	94.551(2)	78.1303(8)
<i>V</i> [Å <sup>3</sup> ]	2353.1(1)	1171.12(4)
<i>D<sub>x</sub></i> [g cm <sup>-3</sup> ]	1.320	1.326
$\mu$ (MoK $\alpha$ ) [mm <sup>-1</sup> ]	0.337	0.339
Scan type	$\phi$ and $\omega$	$\omega$
2 $\theta$ (max) [°]	55	60
Transmission factors (min; max)	0.798; 0.972	0.833; 0.931
Total reflections measured	56591	47462
Symmetry independent reflections	10476	6849
Reflections with <i>I</i> > 2 $\sigma$ ( <i>I</i> )	7818	5150
Reflections used in refinement	10468	6849
Parameters refined	568	284
Final <i>R</i> ( <i>F</i> ) [ <i>I</i> > 2 $\sigma$ ( <i>I</i> ) reflections]	0.0567	0.0416
<i>wR</i> ( <i>F</i> <sup>2</sup> ) (all data)	0.1577	0.1135
Weights: <sup>a)</sup> <i>a</i> ; <i>b</i>	0.0762; 2.0292	0.0552; 0.371
Goodness of fit	1.034	1.040
Secondary extinction coefficient	0.005(1)	—
Final $\Delta_{\max}/\sigma$	0.001	0.001
$\Delta\rho$ (max; min) [e Å <sup>-3</sup> ]	1.42; -0.57	0.31; -0.40

<sup>a)</sup>  $w = [\sigma^2(F_o^2) + (aP)^2 + bP]^{-1}$ , where  $P = (F_o^2 + 2F_c^2) / 3$

## Acknowledgment

We thank the analytical services of the Institute of Organic Chemistry of the University of Zürich for spectra and analyses, and the Swiss National Science Foundation and F. Hoffmann-La Roche AG, Basel, for financial support. G. M. thanks the Polish State Committee for Scientific Research for financial support (Grant No. 3 T09A 04625).

## REFERENCES

1. Huisgen R., *Angew. Chem. Int. Ed. Engl.*, **2**, 565 (1963); Huisgen R., *Angew. Chem. Int. Ed. Engl.*, **2**, 633 (1963).
2. Huisgen R., in '1,3-Dipolar Cycloaddition Chemistry', ed. Padwa, A., Vol. 1, Wiley, New York, 1984, p. 1.
3. Huisgen R., 'The Adventure Playground of Mechanisms and Novel Reactions', Ed. Seeman J.T., American Chemical Society, Washington, D.C., 1994.
4. a) Staudinger H. and Gaule A., *Ber. Dtsch. Chem. Ges.*, **49**, 1897 and 1951 (1916); b) Staudinger H. and Siegwart J., *Helv. Chim. Acta*, **3**, 833 (1920).
5. a) Schönberg A. and Vargha L., *Liebigs Ann. Chem.*, **483**, 176 (1930); b) Schönberg A. and Vargha L., *Ber. Dtsch. Chem. Ges.*, **64**, 1390 (1931); c) Schönberg A., Cernik D. and Urban W., *Ber. Dtsch. Chem. Ges.*, **64**, 2577 (1931); d) Schönberg A., Nickel S. and Cernik D., *Ber. Dtsch. Chem. Ges.*, **65**, 289 (1932).
6. Huisgen R., Fulka C., Kalwinski I., Li, X., Mlostoń G., Rodríguez Moran J. and Pröbstl A., *Bull. Soc. Chim. Belg.*, **93**, 511 (1984).
7. Huisgen R., Langhals E., Mlostoń G., Oshima T. and Rapp J., *Lectures in Heterocycl. Chem.*, **9**, S-1 (1987).
8. a) Huisgen R., Fisera L., Giera H. and Sustmann R., *J. Am. Chem. Soc.*, **117**, 9671 (1995); b) Sustmann R., Sicking W. and Huisgen R., *J. Am. Chem. Soc.*, **117**, 9678 (1995).
9. Huisgen R., Li X., Giera H. and Langhals E., *Helv. Chim. Acta*, **84**, 981 (2001).
10. Huisgen R. and Langhals E., *Tetrahedron Lett.*, **30**, 5369 (1989).
11. Huisgen R. and Li X., *Tetrahedron Lett.*, **24**, 4185 (1983).
12. Mlostoń G. and Heimgartner H., *Polish J. Chem.*, **74**, 1503 (2000).
13. Mlostoń G. and Heimgartner H., in 'The Chemistry of Heterocyclic Compounds', Vol. 59: 'Synthetic Applications of 1,3-Dipolar Cycloaddition Chemistry Toward Heterocycles and Natural Products', Eds. Padwa A. and Pearson W.H., Wiley, New York, 2002, p. 315.
14. Huisgen R., Kalwinski I., Li X. and Mlostoń G., *Eur. J. Org. Chem.*, 1685 (2000).
15. Huisgen R., Li X., Mlostoń G. and Fulka C., *Eur. J. Org. Chem.*, 1695 (2000).
16. Urbaniak K., Mlostoń G., Gulea M., Masson S., Linden A. and Heimgartner H., *Eur. J. Org. Chem.*, 1604 (2005).
17. Mlostoń G., Linden A. and Heimgartner H., *Helv. Chim. Acta*, **74**, 1386 (1991).
18. Huisgen R. and Mlostoń G., *Heterocycles*, **23**, 2201 (1985).
19. Aono M., Terao Y. and Achiwa K., *Chem. Lett.*, 1851 (1987).
20. Huisgen R., Mlostoń G. and Fulka C., *Heterocycles*, **23**, 2207 (1985).
21. Büchel T., Prewo R., Bieri J. H. and Heimgartner H., *Helv. Chim. Acta*, **67**, 534 (1984).
22. a) Kägi M., Linden A., Mlostoń G. and Heimgartner H., *Helv. Chim. Acta*, **79**, 855 (1996); b) Kägi M., Mlostoń G., Linden A. and Heimgartner H., *Helv. Chim. Acta*, **77**, 1299 (1994); c) Petit M., Linden A., Mlostoń G. and Heimgartner H., *Helv. Chim. Acta*, **77**, 1076 (1994); d) Kägi M., Linden A., Heimgartner H. and Mlostoń G., *Helv. Chim. Acta*, **76**, 1715 (1993).
23. a) Gebert A., Linden A., Mlostoń G. and Heimgartner H., *Heterocycles*, **56**, 393 (2002); b) Mlostoń G., Linden A. and Heimgartner H., *Helv. Chim. Acta*, **81**, 558 (1998); c) Mlostoń G., Linden A. and Heimgartner H., *Polish J. Chem.*, **71**, 32 (1997).
24. Meier K.-R., Linden A., Mlostoń G. and Heimgartner H., *Helv. Chim. Acta*, **80**, 1190 (1997).
25. Fisera L., Jaroskova L., Matejkova I. and Heimgartner H., *Heterocycles*, **40**, 271 (1995).
26. a) Hassaneen H.M., Shawali A.S., Fargas D.S. and Ahmed, E.M., *Phosphorus, Sulfur and Silicon*, **113**, 53 (1996); b) Linden A., Awad E.M.A.H. and Heimgartner H., *Acta Crystallogr., Sect. C.*, **55**, 1877 (1999).

27. Awad E.M.A.H. and Heimgartner H., unpublished.
28. Andreasch R. and Zipser A., *Monatsh. Chem.*, **24**, 499 (1903).
29. Johnson C.K., 'ORTEP II', Report ORNL-5138, Oak Ridge National Laboratory, Oak Ridge, Tennessee, 1976.
30. Seyfried M.S., Linden A. and Heimgartner H., in preparation.
31. Malaschichin S., Fu C., Linden A. and Heimgartner H., *Helv. Chim. Acta*, **88**, 3253 (2005).
32. Mlostoń G., Janicka E., Woznicka M., Linden A. and Heimgartner H., *Polish J. Chem.*, **77**, 1805 (2003).
33. Huisgen R., Mlostoń G., Giera H. and Langhals E., *Tetrahedron*, **58**, 507 (2002).
34. a) Polshetiwar V. and Kaushik M. P., *Tetrahedron Lett.*, **45**, 6255 (2004); b) Köster R. and Seidel G., *Z. Naturforsch. B*, **43**, 687 (1988).
35. Greidanus J.W., *Can. J. Chem.*, **48**, 350 (1970).
36. a) Majchrzak A., Janczak, A., Mlostoń G. and Heimgartner H., *Helv. Chim. Acta*, **86**, 2272 (2003); b) Elam E. U. and Davis H. F., *J. Org. Chem.*, **32**, 1562 (1967).
37. Heimgartner H. and Mlostoń G., in 'Electronic Encyclopedia of Reagents in Organic Synthesis', Eds. Paquette L., Rigby J., Crich D. and Wipf P., J. Wiley & Sons, Chichester UK, Article RN00429.
38. Heimgartner H., Mlostoń G. and Romański J., in 'Electronic Encyclopedia of Reagents in Organic Synthesis', Eds. Paquette L., Rigby J., Crich D. and Wipf P., J. Wiley & Sons, Chichester UK, Article RN00504.
39. Hooft R., *KappaCCD Collect Software*, Nonius BV, Delft, The Netherlands, 1999.
40. Otwinowski Z. and Minor W., in 'Methods in Enzymology', Vol. 276, 'Macromolecular Crystallography', Part A, Eds. Carter C.W., Jr. and Sweet R.M., Academic Press, New York, 1997, p. 307.
41. Blessing R.H., *Acta Crystallogr. Sect. A*, **51**, 33 (1995).
42. Altomare A., Casciaro G., Giacovazzo C., Guagliardi A., Burla M.C., Polidori G. and Camalli M., *SIR92, J. Appl. Crystallogr.*, **27**, 435 (1994).
43. Spek A.L., *PLATON, Program for the Analysis of Molecular Geometry*, University of Utrecht, The Netherlands, 2005.
44. Beurskens P.T., Admiraal G., Beurskens G., Bosman W.P., Garcia-Granda S., Gould R.O., Smits J.M.M. and Smykalla C., *PATY: The DIRDIF program system*, Technical Report of the Crystallography Laboratory, University of Nijmegen, The Netherlands, 1992.
45. Beurskens P.T., Admiraal G., Beurskens G., Bosman W.P., de Gelder R., Israel R. and Smits J.M.M., *DIRDIF94: The DIRDIF program system*, Technical Report of the Crystallography Laboratory, University of Nijmegen, The Netherlands, 1994.
46. a) Maslen E.N., Fox A.G. and O'Keefe M.A., in 'International Tables for Crystallography', Ed. Wilson A.J.C., Kluwer Academic Publishers, Dordrecht, 1992, Vol. C, Table 6.1.1.1, p. 477; b) Creagh D.C. and McAuley W.J., *ibid.*, Table 4.2.6.8, p. 219; c) Creagh D.C. and Hubbell J.H., *ibid.*, Table 4.2.4.3, p. 200.
47. Stewart R.F., Davidson E.R. and Simpson W.T., *J. Chem. Phys.*, **42**, 3175 (1965).
48. Ibers J.A. and Hamilton W.C., *Acta Crystallogr.*, **17**, 781 (1964).
49. Sheldrick G.M., *SHELXL97, Program for the Refinement of Crystal Structures*, University of Göttingen, Germany, 1997.